

## RECOMMENDED USAGE RESTRICTIONS FOR PLANT PROTECTION PRODUCTS ON SOUTHERN AFRICAN EXPORT CITRUS

Compiled by:

Vaughan Hattingh, Citrus Research International, tel 021-8828553, fax 021-8828557, [vh@cri.co.za](mailto:vh@cri.co.za)  
and Paul Hardman, Citrus Growers Association of southern Africa, tel 031-7652514, fax 031-7658029, [ph@cga.co.za](mailto:ph@cga.co.za)  
August 2012

The purpose of these restrictions is to ensure compliance with residue tolerances in the countries to which Southern African citrus is exported. The recommendations on container labels are based on the **current registrations** of plant protection products. In terms of the SA Act 36 of 1947 (and equivalent legislation in neighbouring countries) the registration requirements must be adhered to at all times. However, the requirements of importing countries have made it necessary to introduce **further restrictions** in order to comply with maximum residue limits.

The active ingredients of the products are listed alphabetically together with an illustrative brand name. In the case of commodity products which may be sold under different brand names only a single name is shown for convenience. This does not imply endorsement of the particular brand concerned.

Due to the multiple fruit set pattern of lemons, plant protection products may only be used as specified during the first spring **blossoming and fruit set** period. Special caution should be exercised to ensure that pesticides applied to protect later sets do not result in unacceptable residues in fruit remaining on the trees from earlier sets.

The following general statements are applicable to the use of plant protection products on all cultivars:

- \* Growers should ensure that these restrictions are kept handy and are consulted before the application of plant protection products.
- \* The application in accordance with current label requirements will in many instances not ensure that export requirements will be met. The restrictions specified are applicable in addition to the label requirements.
- \* The withholding periods specified on product labels provide an indication of the ability of treatments to conform to South African residue levels. Since overseas requirements are generally more stringent these withholding periods are not adequate unless specifically mentioned in this document.
- \* These restrictions apply to the period during which fruit is present on trees (between blossom and harvest) and not to the period between harvest and the onset of blossom, during which time standard label requirements apply.
- \* All usages apply to normal blossom situations. Under conditions where blossoming occurs over an extended period a more conservative approach must be adopted. Under these conditions treatments should be timed according to the early portion of the blossom.
- \* Particular attention must be given to ensuring that spray machinery is calibrated to apply the correct spray volumes in relation to tree size and that spray operators are trained in the handling and application of plant protection products.
- \* All treatments referred to above must be applied at the registered concentrations.
- \* Alternation of products, where applicable with reference to the restrictions, will reduce the risk of excessive residues of any one chemical and will also reduce selection pressure for resistance.
- \* The additional restrictions in this document do not necessarily provide an indication of the compatibility of the products with integrated pest management and good agricultural practice.
- \* The addition of oil to a treatment, if not registered as such, should be avoided as this may increase the residue level.

All exporting growers should keep accurate spray records so that in the event of exceeding MRLs the reasons can be determined. These records should be retained in safe-keeping for at least 3 years.

**Growers are strongly urged to abide by these restrictions to minimise the risk of residue tolerances being exceeded. However, it must be noted that no absolute guarantee can be given that even by following these guidelines export residue tolerances will in all instances not be exceeded. The efficacy and integrated pest management compatibility of plant protection products listed here are additional considerations that users should bear in mind and which are not covered in this document whatsoever. This document has been compiled with information presently available and in good faith, but with the express condition that the authors, Citrus Research International and Citrus Growers Association of Southern Africa, accept no responsibility whatsoever for any loss or damage resulting directly or indirectly from the use thereof.**

**SUMMARY TABLE OF RECOMMENDED USAGE RESTRICTIONS**

<b>PRODUCT</b>	<b>All markets (including EU) except where other restrictions are specified</b>	<b>CODEX (A) <sup>a</sup></b>	<b>CODEX (B) <sup>b</sup></b>	<b>CANADA</b>	<b>U S A</b>	<b>JAPAN</b>	<b>Other</b>
Acetamidrid/ Mospilan	150d PHI as registered	- Taiwan: 150d PHI as registered for oranges and not later than 90% petal fall for other citrus <sup>d</sup>	-	-	-	-	Korea: 150d PHI as registered for Soft Citrus and not later than 90% petal fall for other citrus.
Aldicarb/ Temik	Not permitted in SA <sup>u</sup> . For other southern African countries: 180d PHI	-	-	-	-	-	-
Amitraz/ Mitac	Not later than 90% petal fall	-	-	150d PHI	-	28d PHI as registered	-
Avermectin/ Agrimec	7d PHI as registered	-	-	-	-	-	-
Azadirachtin	See Pyrethrins	-	-	-	-	-	-
Azinphos-methyl/ Gusathion	Not later than 90% petal fall	120d PHI	120d PHI	21d PHI as registered	Not later than 90% petal fall	Not later than 90% petal fall	Korea: 120d PHI
Azoxystrobin/ Ortiva	77d PHI as registered	-	-	-	-	-	-
Bacillus thuringiensis/ Dipel	0d PHI as registered	-	-	-	-	-	-
Bromopropylate/ Acarol	Not later than 90% petal fall	21d PHI	21d PHI	21d PHI	-	21d PHI	-
Buprofezin/ Applaud	45d PHI as registered	-	-	-	-	-	-
Cadusaphos/ Rugby	0d PHI as registered	-	-	-	-	-	-
Carbendazim (Bavistin, Bendazid, Knowin, Benomyl, Spotless)	120d PHI for oranges and grapefruit and 90d PHI for other citrus	90d PHI for oranges and not later than 90% petal fall for other citrus	14d PHI as registered	14d PHI as registered	14d PHI as registered	14d PHI as registered	Korea: 14d PHI as registered
Chlorfenapyr/ Hunter	Medium cover spray: Before calyx closure (± 3 weeks after petal fall) as registered. Bait spray application (20ml): Not later than mid-December and a 140d PHI as registered	-	-	-	-	-	-
Chlorpyrifos/ Dursban	60d PHI for sprays as registered, 0d PHI for soil and stem applications as registered	-	-	-	-	-	-
Chlorantraniliprole (Rynaxypyr)/ Coragen	7d PHI as registered	-	-	Not later than 90% petal fall	-	Not later than 90% petal fall	Not later than 90% petal fall
Clothianidin/ Dantop	120d PHI as registered	-	-	-	-	-	-

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) <sup>a</sup>	CODEX (B) <sup>b</sup>	CANADA	U S A	JAPAN	Other
Copper	14d PHI as registered	-	-	-	-	-	
Cyhexatin (Azocycloin)/ Sipcatin	Only to be used after harvest and before the onset of blossom	-	-	-	Not permitted	Not permitted	
Cypermethrin	28d PHI as registered	-	-	-	Not later than 90% petal fall	-	
Dichlorprop/ Corasil E	Not later than 90% petal fall	-	150d PHI as registered	-	-	150d PHI as registered	
Dichlorprop-p/ Corasil P	90d PHI as registered	-	-	-	-	-	
Dicofol/ Kelthane	14d PHI (Not later than 90% petal fall) <sup>m</sup>	-	-	-	-	-	Korea: Not later than 90% petal fall for soft citrus and 28d PHI for other citrus.
Difenoconazole/ Score	Not later than 90% petal fall as registered	-	-	-	-	-	
Dimethoate/ Rogor	Not later than 50% petal fall for sprays, Not later than white bud stage for soil applications	-	-	42d PHI as registered	42d PHI as registered	42d PHI as registered	
Dimethyl Didecylammonium Chloride/ Sporekill	Not permitted	-	-	-	-	Not permitted <sup>k</sup>	
Dithiocarbamates (Mancozeb/Maneb)	21d PHI or 28d PHI as registered	Not later than 90% petal fall for Lemons and Grapefruit and 21d or 28d PHI as registered for Soft Citrus and Oranges	-	Not later than end December and only where packhouses have either a non-recycling high pressure spray or non-recycling Deccosol foam curtain and regular (twice daily) cleaning of brushes	Not later than 90% petal fall	Not later than end-January	Korea: 21d PHI or 28d PHI as registered for Soft Citrus and not later than 90% petal fall for other citrus
(E)-8-Dodecen-1-yl acetate + (Z)-8-Dodecen-1-yl acetate/ Checkmate	0d PHI as registered	-	-	-	-	-	

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) <sup>a</sup>	CODEX (B) <sup>b</sup>	CANADA	U S A	JAPAN	Other
Endosulfan/ Thiodan <sup>n</sup>	Not permitted in SA <sup>n</sup> . Not later than 90% petal fall in other southern African countries	- <sup>n</sup>	28d PHI <sup>n</sup>	- <sup>n</sup>	- <sup>n</sup>	28d PHI <sup>n</sup>	- <sup>n</sup>
Ethephon/ Ethrel	Not permitted	-	As registered	As registered	-	Post harvest use not permitted	
Ethoprophos/ MOCAP	0d PHI as registered	-	-	-	-	-	
Etoxazole/ Smite	28d PHI as registered	-	-	-	-	-	
Fenamiphos/ Nemacur	150d PHI as registered	-	-	-	-	-	
Fenazaquin	56d PHI as registered	-	-	-	-	-	
Fenbutatin-oxide/ Torque	7d PHI as registered	-	-	-	-	-	
Fenpropathrin/ Meothrin	28d PHI as registered	185 PHI	-	-	-	-	Korea: 28d PHI as registered for soft citrus and 185d PHI for other citrus
Fenpropathrin + Phenthoate/ Meothrin + Eisan	Not later than 90% petal fall	-	-	-	-	-	
Fenpyroximate/ Mitigate	28d PHI as registered	28d PHI as registered for oranges, 150d PHI and not later than end-October for other citrus	-	150d PHI and not later than end-October for other citrus	-	-	Korea: 28d PHI as registered for soft citrus, 150d PHI and not later than end-October for other citrus
Fipronil/ Regent	Before calyx closure (± 3 weeks after petal fall) as registered	-	-	-	-	-	
Formetanate/ Dicarzol	90d PHI as registered	-	-	-	-	-	
Fosetyl-AI/Aliette	0d PHI as registered	-	-	-	-	-	
Fosthiazate/ Nemathorin	43d PHI as registered	-	-	-	-	-	
Gibberellic Acid	15d PHI as registered	-	-	-	-	-	
Furfural/ Crop Guard	43d PHI as registered	-	-	-	-	-	
Granulovirus (Cryptogran, Cryptex)	0d PHI as registered	-	-	-	-	-	
Guazatine	Post-harvest as registered	Not permitted	-	Not permitted	Not permitted	Not permitted	Korea: not permitted

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) <sup>a</sup>	CODEX (B) <sup>b</sup>	CANADA	U S A	JAPAN	Other
Helicoverpa armigera nucleopolyhedrovirus/ Helicovir	0d PHI as registered	-	-	-	-	-	-
Imazalil (Chloramizol)	Post-harvest as registered	-	-	-	-	-	-
Imidacloprid/ Confidor	212d PHI as registered	-	-	-	-	-	Korea: 212d PHI as registered for Soft Citrus and not later than 90% petal fall for other citrus
Iprodione/ Rovral (Dicarboxamil)	115d PHI as registered for Lemons and Soft Citrus, not later than 90% petal fall for other citrus	Not later than 90% petal fall	115d PHI as registered	-	Not later than 90% petal fall	-	Switzerland and Korea: Not later than 90% petal fall
Isazophos/ Miral	56d PHI as registered	-	-	-	-	-	-
Kresoxim-methyl/ Strobry	Not later than 90% petal fall	56d PHI for Grapefruit & Oranges. Not later than 90% petal fall for Lemons & Soft Citrus	56d PHI	-	-	56d PHI	-
Mercaptothion/ Malathion (fruit fly baiting only)	Only dilute concentration, do not use higher concentrations. 28d PHI <sup>L</sup>	7d PHI	7d PHI	14d PHI	7d PHI	7d PHI	-
Metaxyl M/ Ridomil Gold	30d PHI as registered	-	-	-	-	-	-
Methamidophos/ Citrimet	60d PHI	-	21d PHI as registered	21d PHI as registered	-	21d PHI as registered	Korea: 21d PHI as registered
Methidathion/ Ultracide	No later than 90% petal fall	56d PHI as registered	56d PHI as registered	56d PHI as registered	56d PHI as registered	56d PHI as registered	Korea: Not later than 90% petal fall for oranges, grapefruit & lemons and 56d PHI as registered for soft citrus
Methiocarb/ Mesurol	21d PHI as registered	Not later than the end of January	-	-	Not later than the end of January	Not later than the end of January	Korea: 21d PHI as registered for mandarins and not later than the end of January for others

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) <sup>a</sup>	CODEX (B) <sup>b</sup>	CANADA	U S A	JAPAN	Other
Methomyl (Thiodicarb) / Lannate	60d PHI for all registered usages	28d PHI for all registered usages	28d PHI for all registered usages	28d PHI for all registered usages	28d PHI for all registered usages	28d PHI for all registered usages	
Methyl-parathion / Pennacap	Not later than 50% petal fall	-	-	-	-	-	
Mevinphos	28d PHI	-	3d PHI as registered	3d PHI as registered	-	3d PHI as registered	Korea: 3d PHI as registered
Monocrotophos/ Azodrin	Not for use in SA; 90d PHI in other southern African countries	-	-	-	-	-	
Omethoate/ Folimat	Apply no more than once in a season, not later than beginning of December and ensure at least a 150d PHI	-	-	-	-	-	
Paecilomyces lilacinus/ PL+	0d PHI as registered	-	-	-	-	-	
Parathion/ Parathion	Not later than 50% petal fall	-	-	Not later than 4 weeks after petal fall	-	Not later than 4 weeks after petal fall	
Permethrin/ Last Call	0d PHI as registered	-	-	-	-	-	
Phenthoate/ Eisan	Not later than 50% petal fall	-	-	-	-	-	
Phosphorous acid	0d PHI, 14d PHI or 28d PHI as registered	-	-	-	-	-	
Pirimicarb/ Aphox	14d PHI as registered	-	-	Not later than 90% petal fall	Not later than 90% petal fall	Not later than 90% petal fall	Switzerland: Not later than 90% petal fall Korea: not permitted on lemons
Prochloraz	Post-harvest as registered	-	-	Not permitted	Not permitted	Oranges only	
Profenofos/ Selecron	Between blossom and harvest, use Selecron only once and not more than 100mℓ/100ℓ water at not later than 50% petal fall <sup>Y</sup>	-	-	-	-	-	
Propargite/ Omite	14d PHI as registered	-	-	-	Not later than 90% petal fall for Soft Citrus, “-“ for other citrus	-	
Prothiofos/ Tokuthion	Between blossom and harvest, Prothiofos should be used only once and not later than 90% petal fall	-	-	-	-	-	

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) <sup>a</sup>	CODEX (B) <sup>b</sup>	CANADA	U S A	JAPAN	Other
Pyraclostrobin/ Cabrio	45d PHI as registered	- Taiwan: Not later than 90% petal fall for Soft citrus <sup>d</sup>	-	-	-	-	
Pyrethrin (incl natural Pyrethrum) / Erador	2d PHI as registered	Not later than 90% petal fall	Not later than 90% petal fall	-	-	-	
Pyrimethanil / Philabuster	Post-harvest as registered	Post-harvest as registered. Do not use in HOT water bath. Taiwan: Not permitted for soft citrus	-	-	-	Not permitted	Korea: Not permitted
Pyriproxyfen/ Nemesis	90d PHI as registered	-	-	120d PHI	-	-	Switzerland and Korea: 120d PHI
Sodium ortho-phenyl-phenol	Post-harvest as registered	-	-	-	-	-	
Spinetoram/ Delegate	7d PHI as registered	7d PHI as registered for oranges and 14d PHI for other citrus	-	-	-	-	
Spinosad/ (Tracer/GF120)	1d PHI as registered for fruit fly baiting (GF120), but 28d PHI for other applications (Tracer)	-	-	-	-	-	
Spirodiclofen/ Envidor	14d PHI as registered	-	-	-	-	-	
Spirotetramat/ Movento	60d PHI as registered	-	-	-	-	-	Korea: Not later than 90% petal fall
Tartar emetic/ Tartox	30d PHI as registered	-	-	-	-	-	
Tau-fluvalinate/ Klartan	Not later than mid-November as registered	-	-	-	-	-	
Tebuconazole/ Folicur	Not later than 90% petal fall as registered	-	-	-	-	-	
Teflubenzuron/ Nomolt	Not later than 90% petal fall	-	30d PHI	-	-	30d PHI as registered	Korea: 30d PHI on soft citrus and not later than 90% petal fall for other citrus
Temephos/ Abate	200d PHI	-	-	-	-	-	
Terbufos/ (AC92-100, Counter)	30d PHI as registered and not to be used between December and 1 April	-	-	-	-	-	

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) <sup>a</sup>	CODEX (B) <sup>b</sup>	CANADA	U S A	JAPAN	Other
Tetradifon/ Tedium	Not later than 90% petal fall	-	15d PHI	15d PHI	Not Permitted	15d PHI	Switzerland: Not later than 90% petal fall
Thiabendazole	Post-harvest as registered	-	-	-	-	-	
Thiacloprid/ Calypso	No later than 3 weeks after petal-fall	-	-	-	-	-	
Thiophanate-methyl/ Topsin	Not later than 90% petal fall	-	14d PHI as registered	14d PHI as registered	14d PHI as registered	14d PHI as registered	
Trichlorfon/ Dipterex	10d PHI as registered (28d PHI) <sup>m</sup>	28d PHI	10d PHI as registered	-	28d PHI	-	Korea: 10d PHI as registered for Soft Citrus and 28d PHI for other citrus
Trifloxystrobin/ Flint	76d PHI as registered	-	-	-	-	-	Korea: 76d PHI as registered for soft citrus, otherwise on Valencias only and not later than mid-January
Triflumuron/ Alysistin	Not later than 90% petal fall for lemons and 30d PHI as registered for other citrus	Taiwan: Not later than 90% petal fall for soft citrus and 76d PHI as registered for other citrus <sup>d</sup> Not later than 90% petal fall	30d PHI as registered	60d PHI	Not permitted	Not later than 90% petal fall	Switzerland and Korea: Not later than 90% petal fall
2,4-D	Post-harvest, not more than 250ppm in a packhouse treatment and not more than 250ppm in a pre-degreasing drench	-	-	Post-harvest as registered	Post-harvest as registered	Post-harvest as registered	Korea: not permitted on grapefruit and soft citrus
3,5,6 TPA / Maxim	120d PHI as registered	-	-	-	-	-	

PHI = Pre-harvest interval. - = as for "all markets", that is no additional restrictions apply.



**SUMMARY TABLE OF RESIDUE TOLERANCES**

Chemical	RSA	General export tolerance	Codex	Canada	USA	Japan	Korea
Acetamiprid	0.5	0.5	None	0.5	0.5	2.0	2.0 <sup>13</sup>
Aldicarb	0.2	0.02	0.2	0.1	None <sup>13</sup> , 0.3 <sup>11</sup> (None) <sup>e</sup>	0.2, 0.3 <sup>7</sup>	0.02 <sup>13</sup>
Amitraz	0.2	0.05	0.5 <sup>7</sup>	0.1	None	0.5	0.5 <sup>6</sup> , 0.2 <sup>13</sup>
Avermectin	0.01	0.01	0.01	0.02	0.02	0.01	0.02 <sup>13</sup>
Azadirachtin	None	0.01	None	0.1	Exempt	Exempt	None
Azinphos-methyl	2.0	0.05	1.0	2.0	None	0.01	2.0
Azoxystrobin	0.5	0.5	15.0	0.8	10.0	2.0	1.0 <sup>13</sup> (10.0 <sup>x</sup> )
Bacillus thuringiensis	None	None	None	None	None	None	None
Bromopropylate	3.0	0.01	2.0	2.0	None	2.0	5.0
Buprofezin	0.05	0.05	1.0	0.1	2.5	2.0 <sup>7</sup> , 2.5 <sup>11</sup>	0.3 <sup>7</sup> , 1 <sup>13</sup>
Cadusafos	0.05	0.01	None	0.1	None	0.01	None
Carbendazim	5.0	0.2 <sup>7,19</sup> , 0.7 <sup>13,20</sup>	1.0 <sup>7</sup>	10.0	10.0	7.0 <sup>19,20</sup> , 3.0 <sup>11</sup>	7.0
Chinomethionat	0.5	0.01	None	0.1	None	0.5	0.5
Chlorfenapyr	0.01	0.01	None	0.1	0.01	2.0	1.0 <sup>13</sup>
Chlorantraniliprole	0.5	0.5	0.5	0.1	1.4	0.01	None
Chlorpyrifos	0.3	0.3 <sup>11</sup> , 0.2 <sup>20</sup>	1.0	1.0	1.0	1.0	0.3
Clothianidin	0.01	0.01	None	0.1	0.4	2.0	1.0 <sup>13</sup>
Copper	20.0	20.0	None	50.0	Exempt	Exempt	None
Cyhexatin (Azocytotin)	2.0	0.2	0.2	0.1	Not permitted, 0.1 <sup>18</sup>	Not permitted	2.0
Cypermethrin	0.2	0.2	2.0	1.0	None, 0.35 <sup>c</sup>	2.0	2.0
Dichlorprop-p	None	0.05	None	0.1	None	3.0	None
Dicofol	5.0	2.0 (0.02) <sup>m</sup>	5.0	5.0	6.0	5.0	1.0 <sup>6</sup>
Difenoconazole	0.05	0.05	None	0.1	0.6	0.01	1.0 <sup>13</sup>
Dimethoate	2.0	0.02	5.0	1.5	2.0	2.0	2.0
Dimethyl Didecyl ammonium Chloride	2.0	0.01	None	0.1	None	Not permitted <sup>k</sup>	None
Dithiocarbamate	3.0	3.0	2.0 <sup>7</sup> , 10.0 <sup>13</sup>	0.1	None	2.0 <sup>11</sup> , 10.0 <sup>13</sup>	5.0 <sup>13</sup>
(E)-8-Dodecen-1-yl acetate + (Z)-8-Dodecen-1-yl acetate	None	None	None	None	Exempt	None	None
Endosulfan <sup>h</sup>	1.0 <sup>h</sup>	0.05	None	0.1	None	0.5	0.1 <sup>19,20</sup> , None <sup>7,13</sup>
Ethephon	2.0	0.05	None	1.0	None	2.0	2.0 <sup>20</sup> , 0.5 <sup>13</sup>
Ethoprophos	0.05	0.02	None	0.1	None	0.005	None
Etoxazole	None	0.01	None	0.1	0.1 <sup>6</sup>	0.7	0.5 <sup>13</sup>
Fenamiphos	0.05	0.02	None	0.1	0.6	0.2	0.5 <sup>6</sup>
Fenazaquin	0.05	0.05	None	0.1	0.5 <sup>11</sup> , None <sup>19</sup>	0.01	0.7 <sup>13</sup>
Fenbutatin-oxide (Hexakis)	1.0	1.0	5.0	2.0	20.0	5.0	5.0
Fenpropathrin	0.5	0.5	None	2.0	2.0	5.0	5.0 <sup>13</sup>
Fenpyroximate	0.2	0.2	0.2 <sup>7</sup> , 0.01 <sup>11</sup>	0.1	0.6	1.0	0.5 <sup>13</sup>
Fipronil	0.05	0.005	None	0.1	None	0.01	0.05 <sup>13</sup>
Formetanate	0.5	0.05	None	4.0	1.5 <sup>7,19</sup> , 0.6 <sup>20</sup> , 0.03 <sup>10</sup>	4.0	None
Fosetyl-Al	15.0	15.0	None	9.0	None	150.0	None
Fosthiazate	0.1	0.02	None	0.1	None	0.01	None
Furfural	1.0	1.0	None	None	None	None	None
Gibberellic Acid	0.2	0.2	None	0.1	Exempt	0.2	None

Chemical	RSA	General export tolerance	Codex	Canada	USA	Japan	Korea
Granulovirus	None	None	None	None	None	None	None
Guazatine	5.0	5.0	None	0.1	None	None	None
Helicoverpa armigera nucleopolyhedrovirus	None	None	None	None	None	None	None
Imazalil (Chloramizol)	5.0	5.0	5.0	5.0	10.0	5.0	5.0
Imidacloprid	0.5	0.5	1.0	1.0	0.7	0.7	0.5 <sup>13</sup>
Iprodione	1.0	1.0 <sup>20,10</sup> , 0.02 <sup>11</sup>	None	0.1	None	10.0	None
Isazophos	0.02	0.01	None	0.1	None	0.01	None
Kresoxim-methyl	0.5	0.05	0.5 <sup>22</sup>	0.1	None	10.0	2.0 <sup>13</sup>
Mercaptothion	4.0	0.02	7.0	0.1	8.0	4.0	None
Metalaxyl M (Mefenoxam)	0.5	0.5	5.0	5.0	1.0	0.7	None
Methamidophos	0.2	0.01	None	0.1	None	1.0	0.5
Methidathion	2.0	0.02	2.0	2.0	4.0 <sup>6</sup> , 6.0 <sup>10</sup>	5.0	None <sup>6</sup> , 5.0 <sup>13</sup>
Methiocarb	0.1	0.1	None	0.1	None	0.05	0.05 <sup>6</sup> , 0.5 <sup>13</sup>
Methomyl (Thiodicarb)	0.2	0.01	1.0	1.0	2.0	10.0	1.0
Methyl-parathion	1.0	0.02	None	0.1	None	0.2	None
Mevinphos	0.1	0.01	None	0.2	None	0.2	0.2
Monocrotophos	Not permitted	Not permitted on SA fruit; 0.01 on fruit from other Sthn African countries	None	0.1	None	0.2	0.2
Omethoate	2.0	0.02	None	1.5	None	1.0	0.2 <sup>13</sup> , 0.01 <sup>6</sup>
Parathion	0.5	0.05	None	1.0	None	0.5	None
Permethrin	0.01	0.01	0.5	0.1	None	5.0	0.5
Phenthoate	1.0	0.01	None	0.1	None	0.1	None
Phosphorous acid	50.0	15.0	None	0.1	Exempt	0.01	None
Pirimicarb	0.5	0.5	3.0	0.1	None	0.05	0.05 <sup>11</sup> , 0.5 <sup>2</sup>
Paecilomyces lilacinus	None	None	None	None	None	0.01	None
Prochloraz	2.0	2.0	10.0	0.1	None	10.0 <sup>11</sup> , 5.0 <sup>7</sup>	2.0 <sup>13</sup> , 5.0 <sup>22</sup>
Procymidone	0.2	0.02	None	0.1	None	0.5	None
Profenofos	1.0	0.05	None	0.1	None	0.05	None
Propargite	2.0	2.0	3.0	5.0	5.0 <sup>6</sup>	3.0	5.0
Prothiofos	0.05	0.01	None	0.1	None	0.1	0.2 <sup>13</sup>
Pyraclostrobin	0.5	0.5	1.0	2.0	2.0	1.0	0.5 <sup>13</sup>
Pyrethrins (incl Pyrethrum)	1.0	1.0	0.05	1.0 <sup>2</sup>	Exempt	1.0	1.0
Pyrimethanil	10.0	10.0	7.0	10.0	10.0	None (15.0) <sup>9</sup>	1.0 <sup>13</sup>
Pyriproxyfen	0.2	0.2	0.5	0.1	0.3	0.5	None
SOPP	10.0	5.0	10.0	10.0	10.0	10.0	10.0
Spinetoram	0.05	0.05	0.07 <sup>7</sup>	0.3	0.3	0.3	0.05
Spinosad	0.05	0.05	0.3	0.3	0.3	0.3	0.1 <sup>13</sup>
Spirodiclofen	0.01	0.01	0.4	0.5	0.5	2.0	2.0 <sup>13</sup>
Spirotetramat	0.4	0.4	0.5	0.6	0.6	1.0	None
Tartar emetic (Antimony)	3.0	1.0	None	0.1	None	0.01	None
Tau-Fluvalinate	1.0	0.1	None	0.1	None	2.0	None
Tebuconazole	0.02	0.02	None	0.1	None	5.0	2.0 <sup>13</sup>


Chemical	RSA	General export tolerance	Codex	Canada	USA	Japan	Korea
Teflubenzuron	0.5	0.05	None	0.1	None	1.0	0.7 <sup>13</sup>
Temephos	1.0	0.01	None	0.1	None	0.01	None
Terbufos	0.1	0.01	None	0.1	None	0.005	None
Tetradifon	5.0	0.02	None	2.0	None	3.0	2.0
Thiabendazole	6.0	5.0	7.0	10.0	10.0	10.0	10.0
Thiacloprid	None	0.02	None	0.1	None	0.01	0.3 <sup>13</sup>
Thiophanate-Methyl	5.0	0.01	None	10.0	10.0	7.0 <sup>19,20</sup> , 3.0 <sup>11</sup>	None
Trichlorfon	0.1	0.1 (0.01) <sup>m</sup>	None	0.1	None	0.1	0.1 <sup>13</sup>
Trifloxystrobin	0.1	0.1	0.5	0.1	0.6	0.5	0.5 <sup>13</sup>
Triflumuron	0.5	0.05 <sup>20</sup> 0.5 <sup>11</sup>	None	0.1	None	0.02	0.05
2,4-D	2.0	1.0	1.0	2.0	5.0	2.0	2.0 <sup>7,20</sup>
3,5,6 TPA/Trichlopyr	0.1	0.05	None	0.1	None	0.1	0.1 <sup>13</sup>

"None" = no MRL, therefore fruit must be free of detectable residue



**SUMMARY TABLE OF CHANGES EFFECTIVE FROM THIS EDITION**

<u>Active</u>	<u>Country</u>	<u>Previous MRL</u>	<u>New MRL</u>	<u>PHI Changes</u>
Acetamiprid	Taiwan <sup>d</sup>	0.5	0.01	150d PHI as registered for Oranges and not later than 90% petal fall for other citrus
Aldicarb	USA	None <sup>13</sup> , 0.3 <sup>11</sup>	None <sup>13</sup> , 0.3 <sup>11</sup> (None) <sup>e</sup>	See NOTE u
Azinphos-methyl	Japan	1.0	0.01	Not later than 90% petal fall
Chlorantraniliprole	General Export Tolerance	0.01	0.5	See table changes
Chlorantraniliprole	Codex	0.01	0.5	See table changes
Dicofol	Taiwan <sup>d</sup>	2.0	1.0	28d PHI
Dicofol	General Export Tolerance	2.0	2.0 (0.02) <sup>m</sup>	14d PHI (Not later than 90% petal fall) <sup>m</sup>
Furfural	All	-	See MRL table	See Restrictions table
Imidacloprid	Taiwan <sup>d</sup>	1.0	0.01	212d PHI as registered for Oranges and not later than 90% petal fall for other citrus
Pyraclostrobin	Taiwan <sup>d</sup>	0.5	0.01	Not later than 90% petal fall for Soft citrus
Pyrimethanil	Taiwan <sup>d</sup>	10.0	0.01	Not permitted for Soft Citrus
Trichlorfon	General Export Tolerance	0.1	0.1 (0.01) <sup>m</sup>	10d PHI as registered (28d PHI) <sup>m</sup>
Trifloxystrobin	Taiwan <sup>d</sup>	0.1	0.01	Not later than 90% petal fall for Soft citrus and 76d PHI as registered for other citrus



## NOTES

### Numerical Superscripts:

- 1 fruit without peel/pulp
- 2 just oranges
- 3 exocarp of summer oranges
- 4 fruit (except exocarp of summer oranges)
- 5 just peel
- 6 except mandarin oranges
- 7 Oranges, sweet, sour
- 8 Citrus pulp, dried
- 9 Whole fruit
- 10 Clementines, mandarins
- 11 Other citrus
- 12 Except summer oranges
- 13 Mandarins
- 14 Pulp juice
- 15 No specific crop
- 16 Summer orange and mandarins
- 17 Summer orange, pulp and peel
- 18 Summer orange, pulp
- 19 Only grapefruit
- 20 Only lemons
- 21 mandarins, limes and lemons
- 22 grapefruit, oranges and pommelos.

“None” = no MRL, therefore fruit must be free of detectable residue

### Alphabetical Superscripts:

- a = CODEX (A): See Regions and Country Groupings Table below.
- b = CODEX (B): See Regions and Country Groupings Table below.
- c = USA has a MRL on citrus for Zeta-Cypermethrin of 0.35mg/kg but none for Cypermethrin, which means only Zeta-Cypermethrin plant protection products are suitable for the USA.
- d = Reference to “Citrus Fruits” in Taiwan MRL legislation should be interpreted as “oranges only” but “Citrus” refers to all citrus types.
- e= US Environmental Protection Agency has proposed revoking all citrus tolerances for the nematocide Aldicarb within the next 6 months.
- g = A 15.0 mg/kg MRL will come into effect once Pyrimethanil is approved by the Japanese authorities as a food additive.
- h = From 30<sup>st</sup> April 2012 all Endosulfan uses in South Africa will be prohibited by the Registrar of Act 36 of 1947
- k = Pending decision by Japanese authorities regarding the use of disinfectants.
- L = See CRI Production Guideline for appropriate application techniques.
- m = Draft regulation SANCO/10691/2011 that proposes to reduce the EU MRL for Trichlorfon and Dicofol (all to LOD) was agreed at the March 2012 of the Standing Committee on the Food Chain and Animal Health. It is anticipated these new MRLs will be officially published during the course of the next citrus production and export cycle and therefore it is necessary to advise growers of these anticipated recommended usage restriction changes now.
- u = Aldicarb SA registration has been withdrawn and all aldicarb uses in SA must cease by 31<sup>st</sup> May 2012.
- x = A 10.0mg/kg citrus Azoxystrobin MRL has been proposed although the effective date was not notified (see WTO notice G/SPS/N/TPKM/247).
- y = Pay special attention to the introductory notes on page one (3<sup>rd</sup> paragraph) dealing with on lemons.

### REGIONS AND COUNTRY GROUPINGS

Description	Countries
<b>CODEX A</b> (Where no national MRL is set, CODEX MRLs apply)	<b>Africa</b> [Angola, Benin, Botswana, Congo (Republic of), Gabon, Kenya, Madagascar, Mali, Mauritius*, Mauritania, Namibia, Senegal, Seychelles, Reunion*, Sudan*, Tanzania] <b>Asia</b> [China (Peoples Republic of), Hong Kong, India, Indonesia, Malaysia, Philippines, Singapore, Taiwan, Viet Nam]
<b>CODEX B</b> (Where no national MRL is set, CODEX MRLs apply, or when no CODEX MRL is set RSA MRL apply)	<b>Africa</b> [Burkina Faso, Cameroon, Cote D'ivoire, Malawi, Nigeria, Tunisia, Uganda] <b>Asia</b> [Bangladesh, Sri Lanka] <b>Middle East</b> [Azerbaijan, Bahrain, Iran, Jordan*, Kuwait, Oman, Pakistan, Qatar, Saudi Arabia, , United Arab Emirates] <b>Other</b> [Russian Federation, Georgia*]

\* It has not been possible to re-confirm these requirements recently but presumably they still apply