

# LEARNER GUIDE

## Plant Propagation

### Level 4



P.O. Box 461, Hillcrest, 3650  
(031) 313-3364

<b>Title:</b>	<b>Propagate Plants in a Variety of Situations</b>						
<b>Applied Title:</b>	<b>Propagate Citrus Plants</b>						
<b>Field:</b>	Agriculture and Nature Conservation						
<b>Sub-Field:</b>	Primary Agriculture						
<b>SETA (SGB):</b>	AgriSETA						
<b>Skills Area:</b>	Propagation						
<b>Context:</b>	Citrus Production						
<b>US No:</b>	116316	<b>Level:</b>	4	<b>Credits:</b>	3	<b>Notional Hours:</b>	30
<b>Author:</b>	N. Munduku						

Based on the Production Guidelines of:



Supported by:



**Unit standard alignment and assessment tool development:**

Cabeton Training and Development

**Project coordinator:**

Jacomien de Klerk

**Disclaimer**

By accepting this document and reading its contents you agree to be bound by the terms of this disclaimer.

The use of the contents of this document is at your own risk. Neither the Citrus Academy nor the CRI or the CGA warrant that the content of this document is suitable for your intended use or that it is free of inaccuracies or omissions. The opinions and advice expressed in this document are not necessarily those of the Citrus Academy, the CRI or the CGA. The Citrus Academy, the CRI and the CGA, their directors, officers, employees, agents and contractors shall not be liable for any loss or damage of any nature suffered by any person as a direct or indirect result of the use of, or inability to use any advice, opinion or information contained in this document, or any misrepresentation, misstatement or omission, whether negligent or otherwise, contained in this document.

You indemnify the Citrus Academy, the CRI and the CGA against any claim by any third party against the Citrus Academy, the CRI or the CGA, their directors, officers, employees, agents or contractors arising from, or in connection with, the use of, or reliance on, the contents of this document. It is your responsibility to determine suitability of the contents of this document for your intended use.

## Table of Contents

<b>Directions .....</b>	<b>5</b>
<b>Introduction .....</b>	<b>6</b>
1. Purpose.....	6
2. Learning Assumed to Be in Place.....	6
<b>Revision of Level 3 .....</b>	<b>7</b>
1. Function of Environmental Conditions.....	7
2. Monitoring Environmental Conditions.....	8
3. Propagation Procedures.....	8
4. Tools and Equipment.....	9
<b>Chapter 1.....</b>	<b>11</b>
1. Introduction.....	11
2. Methods of Asexual Propagation .....	11
2.1. Grafting.....	11
2.2. Cuttings.....	12
2.3. Layering .....	13
2.4. Tissue Culture.....	13
3. Asexual Propagation for Other Plants .....	13
4. Use of Hormones for Asexual Propagation .....	14
5. Sanitary Measures in Propagation Procedures .....	14
<b>Chapter 2.....</b>	<b>16</b>
1. Introduction.....	16
2. Propagation Structures .....	16
2.1. Germination Rooms.....	16
2.2. Greenhouses.....	16
2.3. Shade-Houses.....	17
3. Potential Problems with Structures .....	18
4. Propagation Media .....	19
5. Growing Media Mixes .....	20
<b>Chapter 3.....</b>	<b>21</b>
1. Introduction.....	21
2. Role of Artificial Propagation Media .....	21
3. Problems with Propagation Media.....	22
4. Effectiveness of Different Processes .....	22
5. Successful Versus Non-Successful Propagation Media and Environments .....	22
<b>Chapter 4.....</b>	<b>24</b>
1. Introduction.....	24
2. Readiness for Transference to Next Phase .....	24
3. Pests and Diseases.....	24
3.1. Foliar Damaging Pests .....	24
3.1.1. Orange Dog .....	25
3.1.2. Thrips.....	25
3.1.3. Aphids.....	25
3.1.4. Citrus Psylla .....	25
3.1.5. Loopers .....	25
3.1.6. Mites.....	25
3.1.7. Red Scale .....	26
3.1.8. Mealybug.....	26







## Revision of Level 3

### 1. Function of Environmental Conditions

- Environmental conditions impact on the growth rate of plants, by mostly impacting on two metabolic processes, being photosynthesis and respiration, or transpiration.
- During photosynthesis, the plant absorbs CO<sub>2</sub> and water, and uses heat to manufacture sugars (carbohydrates), oxygen, and water molecules.
- During respiration O<sub>2</sub> from the air is used to break down carbohydrates in the plant into energy that is used by the plant, CO<sub>2</sub> that is released into the air, and water molecules that are dispersed from the surface of the leaf.
- When the moisture in the surrounding air increases, plant transpiration decreases and light intensity influences transpiration through leaf surface temperature.
- Three classes of environmental conditions impact on plant growth, being atmospheric conditions, biotic conditions and edaphic conditions.
- Atmospheric conditions refer to light, humidity, temperature, water and aeration.
- Seed germination is activated by increased light intensity and light impacts on the rate of photosynthesis and respiration. Fluorescent lights are used in germination rooms to promote seed germination, and natural light is used in tunnels and shade houses.
- Relative humidity impacts on the transpiration rate of plants.
- Temperature impacts on photosynthesis and respiration rates. Germination requires relatively high temperatures, with a range of between 27°C and 32°C, with an average of 29°C being considered optimal.
- Water plays a role in keeping the plant cells turgid, as a catalyst in biochemical reactions in the plant, and in the translocation of manufactured compounds.
- Aeration in the leaf and root areas allows gaseous exchange that ensures that growth and development take place normally.
- Biotic conditions refer to bacteria, fungi and viruses in the rhizosphere, insects and weeds.
- Certain types of bacteria, fungi and viruses in the root-zone are beneficial and have a positive impact on the growth of the plant, while others are non-beneficial and have a negative impact.
- Insects are also classified as beneficial and non-beneficial. Examples of beneficial insects are *Aphytis lingnanensis*, *Chilocorus nigritus*, and *Cryptolaemas montrouzieri*, while non-beneficial insects include red mites, red scale, mealybugs, aphids, leafminer, and thrips.
- Weeds negatively impact on the growth of seedlings because they compete for water and nutrients.
- Edaphic conditions refer to physical and chemical properties of the growth medium in the root-zone, and the properties of the container in which the plant is grown.
- Soilless media that are also used in nurseries include pine-bark, sawdust, peanut shell, river sand, composted organic material, and ash.





- Budding knives and pruning shears are the main tools that are used during budding.
- Shoot tip grafting is another propagation method that is used to eliminate viruses in the plant material.

#### 4. **Tools and Equipment**

- The tools and equipment used commonly in South African citrus nurseries are seed trays, seedling trays, watering cans, knapsacks, soil augers, jugs, spades, planting containers, mesh panels, hose pipes, stakes, collection bags, wheelbarrows, tying machines (tapeners), rakes, and leather gloves.
- Health and safety is addressed by the Occupational Health and Safety Act (OHSA), which prescribes that conditions that may threaten the health and safety of workers must be anticipated, recognised, evaluated, and controlled.
- Tools and equipment must be stored in clean and dry facility that is kept safe and secure.



<p><b><u>Propagation</u></b></p> <p>Propagation in citrus production refers to the multiplication of plant material that is of a specific cultivar and variety, and that possesses more desirable characteristics, such as yield, fruit size and shape and internal quality.</p>
<p><b><u>Dormancy</u></b></p> <p>Dormancy refers to the ability of certain plant-parts, such as seeds, to suspend metabolic processes until ideal environmental conditions occur.</p>
<p><b><u>Metabolic Processes</u></b></p> <p>Metabolic processes refer to organic chemical processes inside a cell that enable life.</p>
<p><b><u>Humidity</u></b></p> <p>Humidity, also referred to as <i>relative humidity</i>, is the amount of water vapour in the air at a given temperature, and is expressed as a percentage. This means that at 20% relative humidity, 20% of any given volume of air will consist of suspended water molecules.</p>
<p><b><u>Respiration</u></b></p> <p>Respiration refers to the process during which the plant takes up oxygen (O<sub>2</sub>) and releases carbon dioxide (CO<sub>2</sub>).</p>
<p><b><u>Photosynthesis</u></b></p> <p>Photosynthesis refers to the chemical reaction that takes place when the plant takes up CO<sub>2</sub>, which combines with water molecules in the plant to produce carbohydrates, which is food for the plant. O<sub>2</sub> is released during this process.</p>
<p><b><u>Ovule</u></b></p> <p>The ovule is a small structure inside the ovary of a seed plant that contains the female reproductive cells inside the embryo sac, and which develops into a seed after fertilisation.</p>

### **Embryo**

The embryo is a plant in its earliest stage of development before an organism becomes self-supporting. Once the embryo begins to grow out from the seed, or germinate, it is called a seedling.

### **Meiosis**

Meiosis is a type of cell division in organisms that reproduce sexually and results in cells with half the number of chromosomes of the original cell.

### **Rootstock**

Rootstock means the root or part of a root used for plant propagation. In reference to grafting, the rootstock is that part of a grafted plant that supplies the aboveground plant parts.

### **Grafting**

Grafting refers to any process of inserting a part of one plant into or onto another plant in such a way that they will unite and grow as a single unit.

### **Apical Dominance**

Apical dominance refers to powerful tip growth that suppresses the growth of lateral buds.

### **Osmosis**

Osmosis is the flow of water and other liquids through a semi-permeable membrane, such as the thin membrane underneath the shell of an egg, from an area with a low concentration of dissolved matter, such as salts, to an area with a high concentration of dissolved matter, so that the concentration imbalance is gradually evened out.

### **Diffusion**

Diffusion is the spontaneous spreading of something such as particles, heat, or momentum. The phenomenon is readily observed when a drop of coloured water is added to clear water, or when smoke from a chimney dissipates into the air. In these cases, diffusion is the result of turbulent fluid motion rather than chemical reactions or the application of external force. In cell biology, diffusion is described as a form of "passive transport", by which substances cross membranes.

### **Rhizosphere**

The rhizosphere, also referred to as the root-zone, is the soil zone that surrounds the roots of the plant. This zone influences the roots and is in turn influenced by the roots.

### **Edaphic**

The term edaphic means the effect of soil characteristics, especially chemical and physical properties, on plants and animals. For the purpose of this section, edaphic conditions refer to the properties of the growth medium, whether it is soil or another substance.

### **Propagules**

Propagules are the shoots, seeds, or other parts that plants use to spread or propagate, either sexually or vegetatively.





The water holding capacity of the medium must be taken into account. Excessive water supply, or a water-saturated medium, leads to reduced levels of O<sub>2</sub> and reduced root formation. The medium must freely drain excess water. Draining water will take with it most nutrients found in solution.

The condition of the plants must be monitored and corrected with foliar feeds if need be.

With developed roots and active shoots, the cutting grows to a plant that can be transplanted after five to six weeks. The hardening off process consist of gradual reduction of humidity and fog or mist until the plants no longer require moisture on the leaves to prevent wilting.

### 2.3. Layering

Layering is another method that can be used to propagate citrus. Roots are induced to develop from stems while these are still attached to the tree. The rooted stems are then cut and placed in containers or directly into the ground as trees. This method is used mainly on bigger branches. Commercially, this method is not used in the citrus industry.

### 2.4. Tissue Culture

Tissue culture concerns reproducing plants in a sterile and aseptic environment by using portions from a mother plant that has the desired characteristics. Laboratories are the best environments for such operations. The plant part that is to be used for the multiplication is selected and extracted. The reproduction process is initiated by placing the plant part in a prepared medium.

In citrus, this method is used for virus indexing and / or cleaning a plant stock of any known and unknown pathogens, such as the Tristeza virus and the Exocortis bacteria.

A budstick from a candidate plant is disinfected and placed in a medium under sterile conditions with adequate light and temperature to induce bud emergence. The apex containing the meristem is extracted and grafted onto a rootstock grown in dark conditions two weeks before.

Troyer rootstock is for example the most sensitive to Exocortis and will be therefore used for screening the pathogen in the budstick. The same procedure is used for Tristeza virus indexing.

In some citrus producing areas in the world, rootstocks that have a low rate of polyembryony are propagated using this method. One such case is the reported in vitro multiplication of the Chinese mandarin as a rootstock in France. Using this process, the rootstock is propagated commercially on a large scale and sold to producers.



#### Polyembryony

Polyembryony means the formation of more than one embryo in a plant seed.

Despite such reports, the use of this technique for mass propagation of citrus plants is limited to regeneration and proliferation of cultivars that have high economic potential, but are in danger of being lost due to disease or ecological factors.

### 3. Asexual Propagation for Other Plants

Different types of plants are propagated using different methods and different plant sections. Here are a few examples:

- **Strawberry** plants are propagated using its aerial stem, referred to as runner. A section of the runner is cut and rooted to produce another plant.

- Plants of the **lily** family, which includes onions, are propagated using the bulb, which is a shortened stem with thick and fleshy leaves. Bulbs are allowed to generate roots and develop into plants.
- **Bananas** can be propagated by means of offshoots, which are lateral shoots developing from the stem, also referred to as suckers. In commercial banana propagation, tissue culture is used because it provides a much higher yield and better plant uniformity, as well as freedom from diseases.
- **Litchis** are propagated through air layering, which is the regeneration of a vegetative part while still attached to the plant.
- **Grape-vines** are propagated through cuttings, which is a method that differs from layering by the fact that the vegetative part being regenerated is detached from the plant.

Plants in all these categories are also commercially reproduced through tissue culture.

#### **4. Use of Hormones for Asexual Propagation**

Hormones in plants are referred to as **plant growth substances**. These substances are produced by the plant in low concentrations and transported to locations within the plant where they are used. Plant growth substances include auxin, cytokinin, gibberellin, abscisic acid and ethylene, and all play an important role in the physiology of the plant.

Artificial hormones are synthetically produced chemicals that induce similar reactions to the naturally occurring hormones in the plant. Hormone treatments are used in citrus propagation only for the rooting of cuttings. The composition of naturally-occurring auxins responsible for adventitious root initiation was determined, with Indole-3-acetic acid (IAA) identified as the main compound promoting this process.

Two major shortcomings are associated with naturally-occurring IAA. Firstly, the low concentration produced at a time makes it insufficient for rooting cuttings in time. Secondly, because the site of hormone production is not the site of action, it is likely that a cutting might not have sufficient hormones at the right site to induce rooting.

Parallel to the identification of IAA, it was established that synthetic products IBA and Naphthalene acetic acid (NAA) were much more effective in adventitious root initiation and development. In propagation through cuttings, IBA is commonly used, and initiates through a series of bio-chemical reactions the transformation of the callus into the root primordium which will develop into roots.

Other hormones such as cytokinin and gibberellin are used for cell growth and differentiation, and stem elongation respectively. These hormones are produced naturally by the plant in sufficient quantities to carry out these physiological processes.

#### **5. Sanitary Measures in Propagation Procedures**

Sanitation in the propagation of citrus is highly important for the survival of the industry. Any person embarking on this activity has to follow the strict regulations set in place by the Citrus Improvement Programme (CIP).

The first line of defence against contamination is to isolate or strictly control access to the propagation area. Those that have access to this area have to be given clear guidance on how to remain 'clean' and free from any contaminants.

Tools used for propagation purposes should be dedicated to specific areas and operations, and not interchanged. Propagation tools must be disinfected on a regular basis, and stored in a clean, dry area.



## **Chapter 1**

- Asexual propagation implies the use of vegetative plant parts to generate another plant which is true-to-type to the mother plant.
- Citrus is propagated asexually by grafting, cuttings, layering, and tissue culture. The choice of propagation method depends on the expected final product.
- The most commonly used grafting method is the inverted T-cut budding method. Chip budding, using a twig with two buds or more and top-working are other grafting methods that are used for citrus.
- Cuttings refers to when plant material with desirable traits is selected and propagated to become a separate pure-line individual.
- Layering is when roots are induced to develop from stems while these are still attached to the tree.
- Tissue culture concerns reproducing plants in a sterile and aseptic environment (laboratory) by using portions from a mother plant that has the desired characteristics.
- Different types of plants are propagated using different methods and different plant sections.
- Hormones in plants are referred to as plant growth substances and are produced within the plant in low concentrations and transported to other locations within the plant where they are used.
- Citrus propagators have to follow the strict regulations set in place by the Citrus Improvement Programme (CIP).

**citrus  
academy**



Complete activities 1 and 2 in the **Learner Workbook**.





## Chapter 2

After completing this chapter, the learner will be able to:

**Recognise and use propagation structures, facilities and materials under supervision and do independent problem solving in relation to processes**

### 1. Introduction

The propagation of citrus can be seen as a natural process that is undertaken in modified, controlled environments in order to enhance the natural process. These modified environments take into account the immediate surroundings and the requirements for citrus propagation. Structures are established to create environments that can be modified and controlled.

### 2. Propagation Structures

For citrus propagation in South Africa, the Citrus Improvement Programme (CIP) guidelines promote propagation in containers. Container propagation makes use of various structures before the tree is finally planted in the land, including:

- Germination rooms
- Greenhouses
- Shade-houses

#### 2.1. Germination Rooms

Seed trays that have been sown with citrus seeds for the production of rootstocks are placed in germination rooms, where they stay until the seedlings have developed and are ready to be transplanted into individual cavities.

Germination rooms are permanent buildings, simply built with raised walls that are plastered and painted. The standard size for a germination room is 3m x 2m with a ceiling height of 2.5m. A germination room is laid out to retain heat, to allow very little air circulation, and to easily build up and maintain humidity.

Germination rooms are equipped with fluorescent lights as a source of red light, which promote germination.

#### 2.2. Greenhouses

Greenhouses, also referred to as tunnels, are the next structure which the citrus seedlings are taken to in the propagation line. Greenhouses are built in such a way that plant growth and development are promoted and accelerated.

Materials used to build the greenhouse should take into account the environment in which the structure will be built, and the environmental requirements of the plants. If the greenhouse is to be built in a location that is known to be windy, provision must be made to strengthen the structure to withstand wind. If the structure is to be built in conditions that are generally humid, special corrosion-resistant material must be used. At the same time, the design and material used in the greenhouse must contribute to creating the ideal environmental conditions that promote plant growth.



A greenhouse has a metal framework that is built from steel or aluminium that is rust and corrosion resistant. The framework is covered with ultraviolet (UV) resistant polyethylene sheeting. Initially glass was used for covering the metal framework. Now several different plastic materials can be used, but the best results, in terms of lifespan and light penetration, have been obtained with 100 micron UV resistant cross-woven polyethylene.

Greenhouses provide a more favourable environment than the outdoors. The propagator is not at the mercy of nature's extremes in as far as temperature is concerned, and propagation and plant production can take place year-round. With the use of control systems, temperature is controlled to allow normal respiration, even when photosynthesis is reduced, commonly in cloudy weather.

Plants are grown closer together in greenhouses resulting in an increased density per surface area. If water-supply is not a limiting factor, light becomes the most important factor as it relates to photosynthesis. With increased light intensity, coupled with increased relative humidity (rh), comes a reduction in the rate of transpiration as CO<sub>2</sub> diffuses rapidly into the leaves increasing the rate of photosynthesis.

Cooled air ensures that the environment is cooled off and increases light intensity. This reduces the respiration rate resulting in less carbohydrates being used up for respiration and more being available for plant growth. Greenhouses can be equipped to humidify the air when required. Air movement also helps to avoid high temperatures and high levels of CO<sub>2</sub>. The propagator must constantly check the balance between heat build-up and the availability of CO<sub>2</sub> for plant growth.

The following equipment is used for this purpose:

- A wet-wall on one side of the greenhouse, or tunnel, made up of a honeycombed cellulose pad, perforated PVC pipes that run water through the wall, a pump, and a shallow tank.
- A set of extractor or exhaust fans on the other end of the tunnel, that pull air from outside the greenhouse through the wet wall which cools and humidifies the incoming air and at the same time removes the 'old' warm air out of the tunnel.
- An irrigation system and tables or beds where plants are to be placed.
- An automated control system that regulates the temperature, aeration, and light intensity by using ceiling curtains.

Figure 2.1 shows a greenhouse where the wet-wall is at the far end. The extractor fans will therefore be behind the photographer.



**Figure 2.1: Seedlings in a Greenhouse**

### **2.3. Shade-Houses**

Shade-houses are where young citrus trees are kept before being sent to the farmer to be planted in the orchard.





## 5. Growing Media Mixes

Propagators of citrus trees in different parts of the world make use of mixtures to form the final growing medium for the plants to be raised in.

In Australia, for instance, propagators frequently use mixtures of peat and sand constituents for raising trees, for instance 3 parts peat and 1 part sand or 1 part peat and 1 part sand. In other places, such as Egypt, mixes of 1 part sand, 1 part sawdust, 1 part peat, have been used successfully.

Sand is too heavy for use in small cells and other propagation containers. This promotes the use of lightweight materials, such as perlite and vermiculite. The extraction of root systems from cell and plug trays can be difficult where sand is used as a constituent of the mix. This leads to the medium separating from the root system and may cause excessive transplant shock.

For propagation through cuttings, because of the high mist frequency, the ideal mixture or medium is the one that has maximum porosity and minimum WHC.



### Chapter 2

- The propagation of citrus can be seen as a natural process that is undertaken in modified, controlled environments in order to enhance the natural process.
- Container propagation makes use of various structures before the tree is finally planted in the land, including germination rooms, greenhouses, and shade-houses.
- Seed trays that have been sown with citrus seeds for the production of rootstocks are placed in germination rooms.
- Greenhouses are built in such a way that plant growth and development are promoted and accelerated.
- Shade-houses are where young trees are kept before being sent to the farmer to be planted in the orchard.
- Structures are used to provide an environment that is and remains easily controllable.
- Potential faults may occur with the control devices or as a result of the wrong interpretation of information that has been relayed by a machine, or with the structure frame itself.
- Different media are available for citrus propagation, ranging from natural coarse river sand to exfoliated vermiculite.
- The most commonly used media in the South African citrus industry is composted pine bark, followed by coarse river sand.
- Propagators of citrus trees in different parts of the world make use of mixtures to form the final growing medium for the plants to be raised in.



Complete activities 3, 4 and 5 in the **Learner Workbook**.







- Other considerations when choosing a medium is sterilisation, salinity, and freedom from weeds nematodes and pathogens.
- Poor irrigation scheduling can lead to the alteration of the physical properties of the medium.
- The medium must be suitable to the environment where propagation is taking place and to the method that is used.
- Different processes are used to obtain propagation media, such as using organic material and transforming it into growing medium through composting or combining different materials. If done correctly, composting is the best economic option.
- Propagation is considered successful if the medium remains clean and free from contaminants up to the end of the process when the tree is planted in the field.



Complete activities 6, 7 and 8 in the **Learner Workbook**.







- Aphids
- Citrus Psylla
- Looper
- Mites

### **Slow Damaging Pests**

- Red scale
- Mealybug
- Slugs

#### **3.1.1. Orange Dog**

Orange dog is the larva of the citrus swallowtail butterfly. Well-developed larvae have a smooth, green appearance with a brown to black stripe on each side. The larvae feed on new flush and severe damage can be caused as the plant becomes exfoliated.

#### **3.1.2. Thrips**

Thrips are translucent orange-yellow in colour. The insects are wingless at the immature nymphal stages, with adults having two pairs of narrow wings. The damage caused by thrips is found on young tender shoots and leaves that are malformed, resulting in stunted growth. Thrips extract chlorophyll from the infested plant parts, which then become pale.

#### **3.1.3. Aphids**

Winged and wingless black and brown aphids secrete honeydew, which causes leaves to become covered with sooty mould. This in turn reduces the rate of transpiration and photosynthesis of the plant. Leaves malformation has also been associated with aphid damage.

The most worrying damage associated with aphids is its ability to transmit the tristeza virus that causes the tristeza disease in citrus. Trees infected with the disease die in the orchard even as soon as ten years after being planted.

#### **3.1.4. Citrus Psylla**

Citrus psylla nymphs are wingless and yellowish in colour. Adults have large and clear wings, and lay eggs on the edges of young, actively growing leaves. Symptoms of the presence of the pest are white deposits on leaves and deformation of young leaves, caused by the nymphs feeding on them. This damage affects tree vitality to a certain extent.

Citrus psylla however also transmits the bacteria that cause greening disease in citrus. Infected trees grow poorly, produce misshapen, small and sour fruit, and have leaves that are chlorotic.

#### **3.1.5. Loopers**

The citrus looper worm feeds on young and mature leaves of citrus plants, starting from the margins. Immature larvae have a different feeding pattern, first feeding on the upper and lower epidermis and then making holes in the leaves.

#### **3.1.6. Mites**

Red mites are oval-shaped, and feed on leaves and green bark of citrus plants. They prefer the upper surface of leaves, turning them grey, silver or yellow. This affects photosynthesis and transpiration rates.

Silver mites also have an oval-shape, but are flat, straw coloured and translucent. They damage the leaves, which become crinkled and may present with corky brown patches on the under-surface.

### **3.1.7. Red Scale**

Red scale feeds off the sap of leaves, twigs and stem of the plant, reducing the vitality of the plant. The surface area for photosynthesis is affected and little chlorophyll is synthesised. Severe infestations lead to leaf drop and twig die-back.

### **3.1.8. Mealybug**

Citrus mealybug is an oval, pale yellow insect covered by powdery white wax. Adults are slow moving and secrete honeydew and black sooty mould. Heavy infestations cause leaf drop and reduced photosynthesis rate.

### **3.1.9. Slugs**

Slugs occasionally infest citrus seedlings. They mainly eat the leaves of orange and grapefruit trees. Rootstock seedlings occasionally get infested.

## **3.2. Wood Damaging Pests**

Rodents feed on the bark of young citrus trees in the nursery. Severe infestations may lead to the trees drying up. When feeding on smaller seedlings, rodents can cut the stem off.

## **3.3. Root Damaging Pests and Diseases**

The pest that causes root damage is nematodes, which feed on roots causing them to appear darker in colour. Rootlets become stunted, swollen and irregular in shape and appearance. Infested trees cannot tolerate stress and drought conditions.

Diseases of concern are root and collar rot caused by Phytophthora, damping off caused by amongst others Rhizoctonia, CBS (citrus black spot), greening diseases which is a bacterial disease transmitted by Psylla, and Tristeza which is a viral disease transmitted by citrus aphids.

Root and stem rot, as well as damping off, start in the nursery and generally express itself in the nursery before trees are planted in the fields. If the plant is infected at the end of the nursery cycle, the disease might not be detected in the nursery but rather express itself in the field.

Damping off is a seedling disease. It only occurs in the seedling stage of propagation, from germination until after the first transplanting.

Tristeza, black spot, and greening are generally field diseases, meaning that the symptoms appear long after the trees have left the nursery. Even if the disease was contracted in the nursery, there is a great probability that these will only be noticed in the field, unless specific analyses are done before the trees leave the nursery.

## **4. Phases of Propagation**

Citrus is propagated in controlled environments from seed germination to taking out the budded tree for planting in the field. Factors such as light intensity, temperature, humidity and aeration are manipulated to create the most conducive environment for citrus propagation.

Each propagation phase has different environmental requirements and some factors are more important than others during different phases. In the germination phase, light intensity and quality, moisture levels, aeration, and temperature are the most crucial factors. Once the seed has germinated, the seedling requires enough humidity to support optimum transpiration.

Respiration and transpiration rates of the seedling determine the level of carbohydrates in the plant that can be used for growth. The more carbohydrates are used in the transpiration process, the less there is available to the plant for growth. Roots develop to ensure that the plant can sustain itself, absorbing water and nutrients, and anchoring the seedling in the medium.

The seedling development phase is therefore mainly characterised by humidity and light intensity control. Humidity levels, coupled with light intensity, influence the temperature in the structure and around the leaf surface area. High humidity reduces the temperature. The preferred temperature is between 20 and 27°C. Higher temperatures negatively affect the seedling.

During the shade-house phase, the plant is subjected to lower humidity and light intensity levels. The use of thermal sheets in the form of shade cloth to protect the plant from harsher environmental conditions of the immediate outdoor is the most important factor.

Ranging from 20% light reduction to near blackout condition (80%), the propagator selects the level that best suits the prevailing conditions of the area. The plant at this stage is acclimatised to fluctuations of environmental conditions.

From one phase to the next the propagator has to ensure that the transfer is done under optimum environmental conditions to avoid transplant shock.

## **5. Hardening-Off Process**

Before the seedlings are transplanted in the shade-houses, and because of the differences in environmental conditions between the greenhouse and the shade-houses, seedlings are subjected to an acclimatisation process to soften transplant shock.

The seedlings stay for a minimum of a week in a facility where the humidity level is lower and the light intensity brought closer to the ambient surroundings. This leads to an accumulation of carbohydrates, increasing the ability of the plant to withstand its new environment and prevailing conditions.

For cuttings, hardening off begins in the structure where propagation takes place. The humidity level is gradually reduced. The rooted cuttings then become more self-sufficient in carrying out physiological activities such as water and nutrient absorption, photosynthesis and leaf development.

## **6. Possible Problems with Hardening-Off Process**

Disease infested plants will display symptoms of the disease and, because of their reduced ability to survive on their own under harsher conditions, will most probably die. Should they survive, they will grow to become inferior trees. Plants with poorly developed root systems will display their inability to survive under harsher conditions during the hardening off process.

Reducing light intensity might lead to spindly and elongated plants. To correct this, an aluminium coated shade cloth (Aluminet) can be used to diffuse light.

With reduced humidity the plants might be losing too much water and carbohydrates through transpiration. If the situation is not corrected plants start to wilt, the extreme being the death of severely wilted plants.




### **Chapter 4**

- The propagator must have a good sense of timing to match the growth stage to the appropriate environment to achieve the best results.
- For citrus, being a woody perennial plant, transplanting readiness is determined by the

growth stage of the seedling and climatic conditions.

- Foliar damaging pests include fast damaging pests such as orange dog, thrips, aphids, citrus psylla, looper and mites, and slow damaging pests such as red scale, mealybug, and slugs.
- Rodents feed on the bark of young citrus trees in the nursery.
- The pest that causes root damage is nematodes, which feed on roots causing them to appear darker in colour. Diseases of concern are root and collar rot caused by Phytophthora, damping off caused by amongst others Rhizoctonia, CBS (citrus black spot), greening diseases which is a bacterial disease transmitted by Psylla, and tristeza which is a viral disease transmitted by citrus aphids.
- Each propagation phase has different environmental requirements and some factors are more important than others during different phases.
- In the germination phase, light intensity and quality, moisture levels, aeration, and temperature are the most crucial factors. Once the seed has germinated, the seedling requires enough humidity to support optimum transpiration.
- Before the seedlings are transplanted in the shade-houses, seedlings are subjected to an acclimatisation process to soften transplant shock.
- Plants with poorly developed root systems will display their inability to survive under harsher conditions during the hardening off process.

Complete activities 9 and 10 in the **Learner Workbook**.



## Bibliography

### **Publications:**

**Citrus Nurseries and Planting Techniques**, B. Aubert and G. Vullin, 1998, Centre of International Cooperation in Agronomical Research for Development (CIRAD)

**Fundamentals of Horticulture**, J.B. Edmond, T.L. Senn, F.S. Andrews and R.G. Halfacre, 4th edition, 1977, McGraw-Hill

**Growing Media for Ornamental Plants and Turf**, K. Handreck and N. Black, 3rd edition, 2002, University of New South Wales Press Ltd, Sydney

**Plant Propagation: Principles and Practices**, H.T. Hartman, E.D. Kester, F.T. Davies and L.R. Geneve, 6th edition, 1997, Upper Saddle River, N.J., Prentice-Hall

**Guidelines for the Production of Container-Grown Citrus Nursery Trees in South Africa**, A.T.C. Lee, K. Roxburgh, 1993, Outspan Publication

**Crop Pests in Southern Africa: Volume 2 — Citrus and Other Sub-Tropicals**, A.C. Myburgh, 1987, Plant Protection Research Institute

