

Fruit size and crop load prediction for citrus

STEPHAN VERREYNNE

Citrus Research International, Department of Horticultural Science, University of Stellenbosch (sv@cri.co.za)

Summary

Fruit size is a very important factor determining market returns. A lot of controllable and non-controllable factors influence eventual fruit size. Estimating final fruit size and crop load is very useful to plan harvesting and to obtain specific markets for the fruit. By accurately predicting final fruit size at any time of fruit development, one can establish the final fruit size distribution at harvest as well as what the size of fruit on a tree should be at a specific time to meet the export size requirements in order to decide if adjustments to the management programme should be done. Fruit size prediction can be done after the physiological fruit drop period by measuring fruit diameter of a representative sample of 50-150 fruit from 2-5 representative trees bearing a typical crop load or to measure all fruit on one or more representative trees in a block or unit. Fruit diameter of tagged fruit should be measured weekly, bi-weekly or monthly from after physiological fruit drop until harvest to build up a database of long-term mean fruit growth rates to be used for future fruit size prediction. Final fruit size at harvest for each fruit is calculated by adding the predicted growth increment until harvest based on the predicted harvest date to the measured fruit size at a specific date. The fruit size distribution at harvest time is used together with the fruit number per tree to predict the yield or crop load per tree. Fruit number estimation should only be done after the physiological fruit drop period by counting the number of fruit per tree for one or more representative trees or by total fruit removal from one or more representative trees. Yield (crop load) is determined by both the number of fruit per tree as well as the size of the fruit. Based on the percentage fruit in each fruit size category and the mean fruit weight (g) of fruit in each category at harvest, crop load can be predicted. The amount of fruit culled from January until harvest due to the Summer-Fall fruit drop period should be taken into account in the prediction. In South Africa there are many growing areas with distinct climatic conditions which make it difficult to provide an overall or general guide to fruit size prediction.




Fig.1. Citrus fruit growth (Bain, 1958).

Opsomming

Vruggrootte is 'n baie belangrike faktor wat inkomste bepaal. Verskeie beheerbare en nie-beheerbare faktore beïnvloed uiteindelijke vruggrootte. Vruggrootte en oeslading voorspelling is belangrik om oes te beplan en om spesifieke markte vir vrugte te verkry. Akkurate finale vruggrootte voorspelling enige tyd tydens vrugontwikkeling is belangrik om die finale vruggrootte verspreiding by oestyd te bepaal en om te bepaal wat die vruggrootte op 'n spesifieke tyd tydens die seisoen behoort te wees om uitvoer vruggrootte standarde te haal om te bepaal of enige veranderinge aan die bestuursprogram gemaak moet word. Vruggrootte voorspelling kan gedoen word na die fisiologiese vrugvalperiode deur vrugdeursnit van 'n verteenwoordigende monster van 50-150 vrugte vanaf 2-5 verteenwoordigende bome met 'n tipiese oeslading te meet of om al die vrugte op een of meer verteenwoordigende bome in 'n blok of eenheid te meet. Die deursnit van gemerkte vrugte moet weekliks, twee-weekliks of maandeliks vanaf na die fisiologiese vrugvalperiode tot by oestyd gemeet word om 'n databasis van langtermyn gemiddelde vruggroeiementos op te bou wat dan in die toekoms vir vruggrootte voorspelling gebruik kan word. Finale vruggrootte by oestyd vir elke vrug word bepaal deur die voorspelde groei toename tot by oestyd, gebaseer op die voorspelde oesdatum, by te tel by die vrugdeursnit op 'n spesifieke dag. Die vruggrootte verspreiding by oestyd word gebruik saam met die aantal vrugte per boom om die oeslading per boom te bepaal. Die aantal vrugte per boom moet na die fisiologiese vrugvalperiode bepaal word deur die aantal vrugte per boom te tel op een of meer verteenwoordigende bome of om al die vrugte vanaf een of meer bome te verwyder. Oeslading word bepaal deur beide die aantal vrugte per boom en vruggrootte. Oeslading word voorspel deur die % vrugte in elke vruggrootte kategorie en die gemiddelde vruggewig van vrugte in elke kategorie. Die aantal vrugte wat val vanaf Januarie tot oestyd as gevolg van die Somer-Herfs vrugval periode moet ook by die voorspelling ingereken word. In Suid-Afrika is daar baie verbouingsareas met verskillende klimaatstoestande wat dit moeilik maak om algemene riglyne vir vruggrootte voorspelling te gee.

1. Introduction

Fruit size is a very important determining factor for market returns. A lot of factors influence eventual fruit size, such as tree health, nutrition, rootstock, irrigation, quality of the bearing wood, the history of the orchard and seasonal differences due to climate or crop load (Gilfillan, 1987). Citrus fruit growth has been described for Valencia oranges (Bain, 1958; Holtzhausen, 1972) and grapefruit (Button, 1969) and follows a sigmoidal growth curve, with 3 distinct growth stages:

CONTINUED ON PAGE 65

Stage I of cell division, Stage II of cell enlargement and Stage III of fruit maturation (Fig.1). Fruit size is determined by the number of cells as well as cell size. Therefore, optimal climatic conditions during both Stages I and II are very important for potential fruit size at harvest and adverse conditions will have a negative impact on fruit size. Although the majority of fruit growth occurs during Stage II, normally little growth occurs in oranges and grapefruit during Stage III of maturation just before harvest.

2. Why is fruit size prediction necessary?

The objective of fruit size prediction is to accurately predict final fruit size at harvest time early in the season, as early as January, but after the physiological fruit drop period at the start of Stage II of rapid fruit growth, or the cell enlargement stage. There is variation in the size of fruit on a tree at a specific time. Estimating final fruit size and crop load per orchard is very useful to plan harvesting and to obtain specific markets for the fruit due to the fact that some markets are very strict on fruit size requirements.

By accurately predicting final fruit size at any time of fruit development, one can establish:

1. the final fruit size distribution at harvest per tree and therefore per orchard, therefore how much fruit would be in a specific fruit size category. These data can be used to predict final crop load (kg/tree).
2. what the size of fruit on a tree should be at a specific time to meet the export size requirements. If these requirements are not met, certain manipulations and adjustments to the management programme can be done to increase the number of fruit meeting the requirements or changes in the marketing programme may be an option.

Although it is too late in January to apply synthetic auxins for thinning and/or fruit size improvement, hand thinning is still an option. Hand thinning at this stage may result in larger fruit, but the major advantage would be the reduction in the number of fruit to reduce the incidence of alternate bearing if there is a history thereof in the orchard.

Previous work on fruit size prediction was aimed at establishing mathematical models to predict final fruit size and the fruit size distribution of 'Valencia' (Franco and Gravina, 2000; Severino et al., 2004) and 'Navel' (Khurshid and Bevington, 2004) oranges. In contrast to the growth of oranges (Bain, 1958; Holtzhausen, 1972) and grapefruit (Button, 1969) (Fig.1), Koch et al. (1996) established that Satsuma and Clementine mandarins grow linearly after physiological fruit drop, during Stage II, until harvest at 0.33 and 0.26 mm/day on average, respectively, without flattening off of the growth curve during Stage III of maturation. Large fruit remained large and small fruit remained small (Koch et al., 1996).



Fruit size and crop load prediction for citrus

3. Measuring fruit size for fruit size prediction

Fruit size prediction can be done after the physiological fruit drop period by measuring fruit diameter of a representative sample of 50-150 fruit from 2-5 representative trees bearing a typical crop load in order to measure at least 500 fruit per block or growing unit. Measure fruit diameter with a calliper, preferably a digital one. The same tagged fruit should be measured weekly, bi-weekly or monthly from after physiological fruit drop until harvest, to build up a database of long-term mean fruit growth rates to be used for future fruit size prediction. Long-term mean fruit growth rates are useful tools to determine the fruit size increment the last few weeks before harvest. Before physiological fruit drop, fruit growth rates are variable and not all fruit has dropped, making measuring very difficult. If the sample used for fruit size determination is not representative of the block the fruit size predictions would not be accurate. An even more representative way of fruit size measurement is to measure all fruit on one or more representative trees in a block or unit.

The size of these fruit at the measured date is used to predict the fruit size distribution of trees in a block at the time of harvest based on historic (long-term) mean fruit growth rates and the number of fruit can be used to calculate crop load. Final fruit size at harvest for each fruit is calculated by adding the predicted growth increment until harvest based on the predicted harvest date to the measured fruit size at a specific date. The fruit size distribution at harvest time is used together with the fruit number per tree to predict the yield or crop load per tree or per ha or orchard.

4. Fruit number per tree estimation

The number of fruit per tree is another important parameter for yield or crop load prediction. The number of fruit per tree at harvest is determined by both flower intensity and the fruit set achieved. Fruit number estimation should only be done after the physiological fruit drop period. A large proportion of fruit drops after physiological fruit drop until harvest, especially for navel orange trees, called the Summer-Fall fruit drop period. There are different ways to determine the number of fruit per tree:

1. Counting the number of fruit per tree for one or more representative trees per orchard or unit by apply-

CONTINUED ON PAGE 66



ing stickers to, then removing them from fruit. The number of stickers used per tree is counted.

2. Total fruit removal from one or more representative trees after the physiological fruit drop period. These fruit are counted, fruit diameter is determined and the fruit size distribution calculated. Based on the percentage fruit in each size category and the mean fruit weight (g) of fruit in each category at harvest, crop load (kg/tree and ton/ha) can be predicted.

5. Yield or crop load prediction

The prediction of yield or crop load (kg/tree) is essential in managing fruit size. Although a lot of factors influence fruit size, yield (crop load) and fruit number plays a major role. Yield (crop load) is determined by both the number of fruit per tree as well as the size of the fruit. Heavy crops with a larger number of fruit result in smaller fruit and if identified early in the season, fruit number reduction strategies like hand thinning can be done. Not only may crop reduction at this stage result in larger fruit, but may reduce the incidence of alternate bearing. Thirty to 38 % of fruit on navel orange trees can be culled from January until harvest due to the Summer-Fall (S-F) fruit drop period (Lima and Davies, 1984; Lima et al., 1980), but the amount of fruit drop during this period for other cultivars is not well documented. The different factors causing fruit drop in navel oranges were recently reviewed by Moore et al. (2006). Historical fruit drop records for each block are important to estimate the size of the Summer-Fall drop in crop load prediction.

6. Fruit size and crop load prediction under South African conditions

In South Africa there are many growing areas with distinct climatic conditions and it is impossible to provide an overall or general guide to fruit size prediction for all the cultivars in all the areas. Using historical long-term means of weekly, bi-weekly or monthly fruit diameter measurements for specific orchards or units, mean daily, weekly, bi-weekly or monthly growth rates until harvest can be calculated. The assumptions are made that small and large fruit have the same growth gradient, for a specific season, therefore a mean fruit growth rate is used for all fruit size categories. These long-term mean growth rates are merely guidelines and the refinement and implementation of fruit size and crop load prediction should be done on the farm itself. Although these mean growth rates are fairly

stable, it may differ with seasonal differences in crop load and climate, humidity, moisture stress etc. Fruit from individual orchards may also differ in growth rate, due to differences in tree age, rootstock, nutrition, irrigation, crop load and tree health (Gilfillan, 1987). Not only would the growth gradient be lower or higher due to such factors, but the growth curve may flatten off earlier or later as well, even for mandarins, in contrast to the observations by Koch et al. (1996).

For example, a representative Washington navel orange tree in Citrusdal at a spacing of 5 x 3 m (667 trees/ha) has 500 fruit (counted by one of the two methods mentioned in section 4) on 15 January. Therefore, all fruit will grow approximately 26.35 mm until harvest according to the fruit growth increments in Table 1 for Citrusdal and the predicted harvest date, say end of June, based on historic data of 3 consecutive years. Yield or crop load, tree health or full bloom date was not included in the calculations. To get the final fruit size at harvest, the growth increment is added to the measured fruit size. Therefore, minimum diameter on 15 Jan (column 4, Table 2) is determined by subtracting 26.35 mm from the minimum diameter of each fruit size category (column 2). Only 100 fruit were measured in this case (column 5), but all the fruit on a tree would be more representative. Fruit diameter of the 100 fruit measured are sorted into the different size categories (column 5). Based on the total number of fruit per tree (500) and the representative fruit size distribution on 15 Jan. (columns 5 and 6), the kg per tree in each category at harvest time can be determined. These calculations can easily be done on an Excel spreadsheet. Assuming 30 % (Lima and Davies, 1984; Lima et al., 1980) of fruit are culled from January until harvest due to the Summer-Fall fruit drop period, only 350 fruit out of the 500 will be harvested. Therefore, the historic cull % during this time for a specific orchard should be documented and subtracted, assuming the same % will be culled from each fruit size category. Mean fruit weight for each fruit size category (column 3) is multiplied by the number of fruit in each size category after subtracting the Summer-Fall drop fruit (column 7) to obtain the kg fruit in each fruit size category (column 8). Therefore, yield or crop load in this example would be 74.8 kg per tree and 49.9 ton/ha.

Growers are encouraged to send their fruit growth measurements to Stephan Verreyne at sv@cri.co.za in order to build up a database of long-term fruit growth data for different cultivars in different climatic areas throughout the country, for more accurate fruit size prediction.

References cited

- BAIN, J.M. 1958. Morphological and physiological changes in the developing fruit of the 'Valencia' orange [*Citrus sinensis* (L.) Osbeck]. Aust. J. Bot. 6:1-24.
- BUTTON, J. 1969. Fruit development in *Citrus paradisi* (Macf. W. Marsh)

seedless in five localities with special reference to the rind. Thesis. Dept. of Horticulture, University of Natal, Pietermaritzburg, South Africa.

FRANCO, J.E. AND GRAVINA, A. 2000. Early prediction of fruit size at harvest for 'Valencia' orange using experimental data. Proc. Intl. Soc. Citricult. 1:456-458.

GILFILLAN, I.M. 1987. Factors affecting fruit size in 'Tomango' and 'Valencia' oranges and practical measures for its improvement. Citrus J. 638:7-13.

HOLTZHAUSEN, L.C. 1972. 'n Morfologiese en fenologiese studie van die blom en vrug van *Citrus sinensis* (L.) Osbeck, cultivar Valencia. Thesis. Dept. of Horticulture, University of Pretoria, South Africa.

KHURSHID, T. AND BEVINGTON, K.B. 2004. Prediction of final fruit size at early stages of fruit development for 'Washington' Navel oranges in Australia with cubic smoothing spline models. Proc. Intl. Soc. Citricult. 2:484-487.

KOCH, N., THERON, K. AND RABE, E. 1996. Fruit size prediction in 'Clementine' and 'Satsuma' mandarin and improvement of 'Clementine' mandarin (*Citrus reticulata* Blanco) fruit size by 2,4-DP. Proc. Intl. Soc. Citricult. 2:737-744.

LIMA, J.E.O. AND DAVIES, F.S. 1984. Growth regulators, fruit drop, yield, and quality of Navel orange in Florida. J. Amer. Soc. Hort. Sci. 109:81-84.

Lima, J.E.O., Davies, F.S. and Krezdorn, A.H. 1980. Factors associated with excessive fruit drop of Navel Orange. J. Amer. Soc. Hort. Sci. 105:902-906.

MOORE, S.D., SCHUTTE, G.C. AND BARRY, G.H. 2006. Factors causing fruit drop in navel oranges. SA Fruit J. 4(6):34-38.

SEVERINO, V., GRAVINA, A. AND FRANCO, J.E. 2004. Early prediction of fruit diameter, and percentage of fruits reaching the commercial size at harvest. Proc. Intl. Soc. Citricult. 2:539-542.

MM FRUIT DIAMETER PER MONTH									
	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
Mihowase Satsuma	9.5	10.1	7.4	8.2	2.6				
Nules Clementine		5.0	7.5	7.6	5.5	3.3			
Eureka Lemon		4.4	5.2	7.2	5.5	3.4	2.4	1.6	
Navel (mean)		4.8	7.3	7.8	4.5	3.9	1.5	0.5	
Bahianinha		6.1	7.0	8.0	4.0	3.7	1.4		
Palmer		3.1	7.8	8.3	5.4	2.9	1.6		
Robyn		5.3	6.6	7.5	3.9	5.3	1.7	0.7	
Washington		3.9	7.8	7.4	4.7	3.8	0.7		
Valencia (mean)		4.5	6.0	7.6	5.4	4.3	2.8	1.7	0.9
Delta		4.4	5.7	7.6	5.1	4.4	3.0	1.4	0.9
Midknight		5.3	6.3	7.8	6.2	4.4	2.4	1.7	0.6
Valencia		3.8	6.2	7.6	4.8	4.3	2.9	1.9	1.2

Table 1. Monthly fruit growth increments in Citrusdal based on historical data from 3 consecutive years. Navel and Valencia in grey represent the means of all Navel or Valencia cultivars, respectively.

Fruit/ 15 kg carton	Minimum diameter (mm)	Mean fruit weight (g)	Minimum diameter on 15 Jan.	% by number on tree 15 Jan.	No. of fruit in category/ tree	After Summer- Fall drop	Kg per tree
32	99	468	72.65	0	0	0	0
36	95	417	68.65	0	0	0	0
40	90	375	63.65	0	0	0	0
48	86	313	59.65	5	25	17.5	5.5
56	82	267	55.65	15	75	52.5	14.0
64	78	234	51.65	30	150	105	24.6
72	73	208	46.65	20	100	70	14.6
88	69	171	42.65	15	75	52.5	9.0
105	65	143	38.65	10	50	35	5.0
125	62	120	35.65	5	25	17.5	2.1
		15 kg/no. of fruit /carton	Min. diameter minus 26.35 mm	out of 100 fruit measured	out of 500 per tree	minus 30% of fruit	Fruit weight x no. after S-F drop

Table 2. Fruit size categories for oranges. Representative fruit size measurements are used to determine fruit size distribution and kg/tree.